PARAGON INFUSION CENTERS

RHEUMATOLOGY ORDER FORM

P: 877.365.5566 | F: 855.889.2946

PATIENT INFORMATION

☐ *Demographics attached*

Patient Name: _____ DOB: _____ Phone: _____

INSURANCE INFORMATION: PLEASE ATTACH COPY OF PRESCRIPTION/MEDICAL CARD(S) (FRONT AND BACK)

MEDICAL INFORMATION

Patient Weight: _____ lbs. Patient Height: _____

Allergies: _____

Previously Failed Therapies: _____ Date: _____

Diagnosis: _____ **(ICD-10)**

- ☐ Iridocyclitic (Uveitis), Unspecified Acute and Subacute
- ☐ Unspecified Iridocyclitis
- ☐ Arthropathic Psoriasis, Unspecified
- ☐ Rheumatoid Arthritis with Rheumatoid Factor, Unspecified
- ☐ Rheumatoid Arthritis without Rheumatoid Factor, Unspecified
- ☐ Rheumatoid Arthritis, Unspecified
- ☐ Ankylosing Spondylitis, Unspecified
- ☐ Gout
- ☐ Systemic Lupus Erythematosus
- ☐ Other: _____

Premedication (if applicable)

- ☐ Tylenol 1000mg PO
- ☐ Benadryl 25-50mg PO/IV
- ☐ Solu-Medrol 125mg IV
- ☐ Pepcid 20mg IV
- ☐ Loratadine 10mg PO **OR** Cetirizine 10mg PO
- ☐ Other: _____

☐ Clinical/Progress Notes, Labs, and Tests supporting primary diagnosis attached

Lab Orders: _____

RHEUMATOLOGY ORDERS

DRUG	DOSING	REFILL
Actemra	☐ 4mg/kg IV every 4 weeks for _____ doses, then followed by 8mg/kg every 4 weeks thereafter ☐ 4mg/kg IV evey 4 weeks ☐ 8mg/kg IV every 4 weeks ☐ Other dose: _____mg IV every 4 weeks	
Cimzia	Initial Dose: 400mg subcutaneously at weeks 0, 2, and 4 weeks Maintenance Dose: ☐ 200mg subcutaneously every 2 weeks ☐ 400mg subcutaneously every 4 weeks	
Krystexxa	☐ 8mg IV in 250mL of NS IV over 120 minutes every 2 weeks *Patient will be observed 1 hour post infusion * Serum Uric Acid required within 72 hours of infusion*	
IVIG	☐ IV ☐ Sub Q ☐ 5% ☐ 10% ☐ _____ % _____ gm/kg IV divided over _____ day(s) _____ mg/kg IV divided over _____ day(s) ☐ Every _____ weeks for one year or _____ one time dose	
Orencia	Orencia Dose: _____mg IV Frequency: ☐ Every 4 weeks OR ☐ 0, 2, 4, weeks, and every 4 weeks thereafter	
SimponiARIA	☐ Initial Dose: 2mg/kg at weeks 0, 4, and then every 8 weeks ☐ Maintenance Dose: 2mg/kg every 8 weeks	
Stelara	☐ Initial Dose: 45mg subcutaneously initially, 4 weeks later, followed by 45mg every 12 weeks ☐ Maintenance Dose: 45mg subcutaneously every 12 weeks ☐ Initial Dose: 90mg subcutaneously initially, 4 weeks later, followed by 90mg every 12 weeks ☐ Maintenance Dose: 90mg subcutaneously every 12 weeks	
Remicade	Dose: _____mg/kg Frequency: ☐ Every _____ weeks ☐ 0, 2, 6, then every 8 weeks	
Rituxan	Dose: 1000mg Frequency: ☐ One time dose ☐ Day 0, repeat dose in 2 weeks	

PHYSICIAN INFORMATION

By signing this form and utilizing our services, you are authorizing *Paragon Healthcare, Inc.* and its employees to serve as your prior authorization and specialty pharmacy designated agent in dealing with medical and prescription insurance companies.

Physician Signature: _____ Physician Name: _____ Date: _____

Phone: _____ Fax: _____ Contact Person: _____

INFUSION CENTER LOCATIONS

☐ Alpharetta, GA ☐ Arlington, TX ☐ Atlanta, GA ☐ Austin, TX ☐ Bee Cave, TX ☐ Birmingham, AL ☐ Cape Coral, FL ☐ Clear Lake, TX ☐ Clearwater, FL ☐ Coppell, TX ☐ Creve Coeur, MO ☐ Dallas, TX ☐ Decatur, GA ☐ Denver, CO ☐ El Paso East, TX ☐ El Paso West, TX ☐ Fort Myers, FL ☐ Fort Worth, TX ☐ Hendersonville, TN ☐ Houston, TX ☐ Huntsville, AL ☐ Independence, MI ☐ Kansas City, MI ☐ Knoxville, TN ☐ Kyle, TX ☐ Lubbock, TX ☐ Montgomery, AL ☐ Murfreesboro, TN ☐ Naples, FL ☐ Nashville, TN ☐ North Hills, TX ☐ Plano, TX ☐ Round Rock, TX ☐ San Antonio, TX ☐ Sarasota, FL ☐ Smyrna, GA ☐ St. Louis, MI ☐ Stone Oak, TX ☐ Waco, TX ☐ West Houston, TX ☐ The Woodlands, TX ☐ Other: _____

IMPORTANT NOTICE: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, privileged property, or exempt from disclosure under applicable law. If you are not the named addressee, you should not disseminate, distribute, or copy this fax. Please notify the sender immediately. If you have received this in error, destroy the document immediately.

© Paragon Healthcare, Inc. 2022