

# IMMUNOGLOBULIN (IG) IV AND SUBQ ORDERS

**P:** 877.365.5566 | **F:** 855.889.2946

PATIENT IN	FORMATI	ON:	Fax completed form, insurance	e information, and clir	ical documentation to 855.889.2946
Patient Name: _					Phone:
Patient Status:	☐ New to 7	Therapy □	Continuing Therapy <b>D</b>	ate of last infusion	on:
MEDICAL INFORMATION					
ICD-10 Code (required):			ICD-10 description:		
Patient Wt:	kg Height:		_ Diabetic □ Yes □ No If obese, use adjusted body wt? □ Yes □ No		
Allergies:		Brand previously used:			
THERAPY ORDER					
□ IV □ SubQ Pharmacist to identify clinically appropriate brand/infusion rates. May substitute based on product availability.					
Loading Dose (as applicable)	☐ mg/kg ☐ gm/kg		x day(s) <b>OR</b> divided over day(s)		☐ One time dose
					☐ Other:
		grams	x day(s) <b>OR</b> divided	l over day(s)	* Give maintenance dose
		_ grains			weeks after loading dose*
Maintenance Dose		□ mg/kg			(s) Q weeks x1 year
		□ gm/kg	x day(s) <b>OR</b> divided	dover day(s)	
		grams	X day(3) <b>GR</b> divided	. over day(3)	
□ Do not substi	tuta Admini				
<ul> <li>Do not substitute. Administer brand:</li> <li>Infuse entire contents of Ig infusion bag/vial(s) per current dose.</li> </ul>					
<ul> <li>If needed, round dose to nearest whole 5 gm vial for IV doses and nearest single-use vial size for subQ doses.</li> </ul>					
Pre-Medication Orders: to be administered 15-30 minutes before infusion					
			□ Normal Saline 500mL IV □ Cetirizine 10mg PO		
			□ Diphenhydramine 25mg PO □ Quzyttir 10mg IVP		
☐ Loratadine 10mg PO ☐ Diphenhydramine 25mg IV ☐ Other:					
Lab Orders: Lab frequency:   Each infusion   Other:					
Required labs to be drawn by 🗆 Paragon Healthcare 🗆 Referring Provider					
Anaphylactic Reaction Orders:  • Epinephrine (based on patient weight)					
• >30kg (>66lbs): EpiPen® 0.3mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x 1					
<ul> <li>15-30kg (33-66lbs): EpiPen® 0.15mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x 1</li> <li>Diphenhydramine - Administer 25-50mg orally OR IV (adult), refer to provider orders or policy for pediatric dose</li> </ul>					
NS 500 mL IV bolus as needed for IVIg therapy (adult), refer to provider orders or policy for pediatric bolus					
Flush orders: NS 1-20mL pre/post infusion PRN and Heparin 10U/mL or 100U/mL per protocol as indicated PRN  *FOR PARAGON USE ONLY					
		_			
Drug/Brand Sele			ND/Dbarmania	st Signature:	Date:
NP/Pharmacist	Name:			st Signature.	
PROVIDER INFORMATION					
agent in dealing with med	ical and prescription	insurance compa	nies, and to select the preferred site of ca	re for the patient.	uthorization and specialty pharmacy designated
Provider Name:			Signature:		Date:
Provider NPI:		_ Phone: _	Fax:	Contac	Date: ct Person:
Upt out of Paragon selecting site of care (if checked, please list site of care):					
PREFERRED LOCATION					
					9%30
City:		State: _		View our	locations here:
					<b>◎ 🕰 🚟</b>



# COMPREHENSIVE SUPPORT FOR IMMUNOGLOBULIN THERAPIES

### REQUIRED DOCUMENTATION FOR INSURANCE APPROVAL

**GENERAL REQUIREMENTS** 

- Patient demographics
- Insurance information
- All applicable diagnoses
- · History and physicial
- Recent progress notes within 12 months
- Patient's height and weight
- Drug allergies
- Physician Orders
- Plus one of the following

### COMMON VARIABLE IMMUNODEFICIENCY (CVID) / HYPOGAMMAGLOBULINEMIA / PARKINSON'S DISEASE (PD)

- Lab last showing Ig levels and subclasses Ig levels.
- Documentation of recurrent infections
- History of antibiotic usage showing failure to respond to antibiotics
- Documented inadequate response to pneumococcal vaccine or tetanus/diphtheria

## CHRONIC INFLAMMATORY DEMYELINATING POLYNEUROPATHY (CIDP) / GUILLAIN-BARRÉ SYNDROME (GBS)

- Labs
- Nerve conduction study, electromyography (EMG)
- Nerve and/or muscle biopsy (if available)
- Nerve conduction velocity (NCV) test results
- Tried and failed treatments
- Spinal tap (if available)

#### MYASTHENIA GRAVIS

- Exacerbation
- Any history of crisis
- Thymectomy

- Any symptoms that affect respiration, speech or motor function
- Tried and failed treatments