



A Carelon Company

IMMUNOGLOBULIN (IG)
IV AND SUBQ ORDERS

P: 877.365.5566 | F: 855.889.2946

PATIENT INFORMATION:

Fax completed form, insurance information, and clinical documentation to 855.889.2946

Patient Name: _____ DOB: _____ Phone: _____

Patient Status: [] New to Therapy [] Continuing Therapy Date of last infusion: _____

MEDICAL INFORMATION

ICD-10 Code (required): _____ ICD-10 description: _____

Patient Wt: _____ kg Height: _____ Diabetic [] Yes [] No If obese, use adjusted body wt? [] Yes [] No

Allergies: _____ Brand previously used: _____

THERAPY ORDER

[] IV [] SubQ Pharmacist to identify clinically appropriate brand/infusion rates. May substitute based on product availability.

Table for Loading Dose (as applicable) with columns for dose units (mg/kg, gm/kg, grams), frequency (x ___ day(s) OR divided over ___ day(s)), and administration instructions (One time dose, Other: _____, * Give maintenance dose ___ weeks after loading dose*).

Table for Maintenance Dose with columns for dose units (mg/kg, gm/kg, grams), frequency (x ___ day(s) OR divided over ___ day(s)), and administration instructions (Q _____ weeks x1 year, Other: _____).

[] Do not substitute. Administer brand: _____

- Infuse entire contents of Ig infusion bag/vial(s) per current dose.
• If needed, round dose to nearest whole 5 gm vial for IV doses and nearest single-use vial size for subQ doses.

Pre-Medication Orders: to be administered 15-30 minutes before infusion

- [] Acetaminophen 500mg PO [] Normal Saline 500mL IV [] Cetirizine 10mg PO
[] Solu-Medrol _____ mg IVP [] Diphenhydramine 25mg PO [] Cetirizine 10mg IVP
[] Loratadine 10mg PO [] Diphenhydramine 25mg IV [] Other: _____

Lab Orders: _____ Lab frequency: [] Each infusion [] Other: _____

Required labs to be drawn by [] Paragon Healthcare [] Referring Provider

Home IV Biologic Ana-kit Orders (adult):

- Epinephrine: >30kg (>66lbs): EpiPen 0.3mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1
• Diphenhydramine: Administer 25-50mg orally OR IV (adult)
• NS 0.9% 1000mL IV bolus per protocol PRN (adult)

Home biologic injection Ana-kit (adult):

- Dispense per protocol EpiPen 0.3mg IM (2-pack)

Refer to physician order or institutional protocol for pediatric dosing Ana-kit

Flush orders: NS 1-20mL pre/post infusion PRN and Heparin 10U/mL or 100U/mL per protocol as indicated PRN

Supply IV Infusion Pump (E0781) and/or SubQ Infusion Pump (E0779) as needed

*FOR PARAGON USE ONLY

Drug/Brand Selection: _____ Date: _____

NP/Pharmacist Name: _____ NP/Pharmacist Signature _____

PROVIDER INFORMATION

By signing this form and utilizing our services, you are authorizing Paragon Healthcare, Inc. and its employees to serve as your prior authorization and specialty pharmacy designated agent in dealing with medical and prescription insurance companies, and to select the preferred site of care for the patient.

Provider Name: _____ Signature: _____ Date: _____

Provider NPI: _____ Phone: _____ Fax: _____ Contact Person: _____

[] Opt out of Paragon selecting site of care (if checked, please list site of care): _____

PREFERRED LOCATION

City: _____ State: _____

View our locations here:



PARAGONHEALTHCARE.COM

IMPORTANT NOTICE: This fax is intended to be delivered only to the named address and contains material that is confidential, privileged property, or exempt from disclosure under applicable law. If you are not the named addressee, you should not disseminate, distribute, or copy this fax. Please notify the sender immediately and destroy all copies if you have received this document in error.



PHI-REF-ORD-10021-V8

REQUIRED DOCUMENTATION FOR INSURANCE APPROVAL
GENERAL REQUIREMENTS

- Patient demographics
- Insurance information
- All applicable diagnoses
- History and physical
- Recent progress notes within 12 months
- Patient's height and weight
- Drug allergies
- Physician Orders
- Plus one of the following

**COMMON VARIABLE IMMUNODEFICIENCY (CVID) /
HYPOGAMMAGLOBULINEMIA / PARKINSON'S DISEASE (PD)**

- Lab last showing Ig levels and subclasses Ig levels.
- Documentation of recurrent infections
- History of antibiotic usage - showing failure to respond to antibiotics
- Documented inadequate response to pneumococcal vaccine or tetanus/diphtheria

**CHRONIC INFLAMMATORY DEMYELINATING POLYNEUROPATHY (CIDP) /
GUILLAIN-BARRÉ SYNDROME (GBS)**

- Labs
- Nerve conduction study, electromyography (EMG)
- Nerve and/or muscle biopsy (if available)
- Nerve conduction velocity (NCV) test results
- Tried and failed treatments
- Spinal tap (if available)

MYASTHENIA GRAVIS

- Exacerbation
- Any history of crisis
- Thymectomy
- Any symptoms that affect respiration, speech or motor function
- Tried and failed treatments